§51.3200

than one-fifth of this tolerance, or 2 percent, may be allowed for onions which are affected by decay or wet sunscald.

(2) U.S. Combination grade. When applying the foregoing tolerances to this grade no part of any tolerance shall be allowed to reduce, for the lot as a whole, the 50 percent of onions of the U.S. No. 1 grade, but individual containers shall have not less than 40 percent of the U.S. No. 1 grade.

(b) For size:

- (1) Not more than 5 percent of the onions in a lot may be smaller than the minimum diameter specified. In addition, not more than 10 percent of the onions in a lot may be larger than the maximum diameter specified.
- (2) When a percentage of the onions is specified to be a certain size and larger, individual packages containing more than 10 pounds may have not less than one-half of the percentage specified: *Provided*, that the entire lot averages within the percentage specified.

SAMPLES FOR GRADE AND SIZE DETERMINATION

§51.3200 Samples for grade and size determination.

Individual samples shall consist of at least 20 pounds for onions packed to meet larger than 21/4 inches maximum diameter. When individual packages contain 20 pounds or more and the onions are packed for Large or Jumbo size or larger the package shall be the sample. When individual packages contain less than 20 pounds, a sufficient number of adjoining packages are opened to provide at least a 20 pound sample, except that for onions packed to meet 21/4 inches maximum diameter or smaller, the sample may consist of either 20 pounds or the individual package.

APPLICATION OF TOLERANCES

§ 51.3201 Application of tolerances.

Individual samples are subject to the following limitations:

(a) Samples which contain more than 20 pounds shall have not more than one and one half times a specified tolerance of 10 percent or more, and not more than double a specified tolerance of less than 10 percent, except that at

least one defective and one off-size onion may be permitted in any sample: *Provided*, that en-route or at destination when onions in containers of 50 pounds or more are packed to a minimum size of 3 inches or larger not more than three onions or more than 4 percent (whichever is the larger amount) may be affected by decay or wet sunscald: *And provided further*, that the averages for the entire lot are within the tolerances specified for the grade; and,

(b) Samples which contain 20 pounds or less shall have not more than double the tolerance specified, except that at least one defective and one off-size onion may be permitted in any sample: *Provided*, that the averages for the entire lot are within the tolerances specified for the grade.

DEFINITIONS

§ 51.3202 Similar varietal characteristics.

Similar varietal characteristics means that the onions in any container are similar in color, shape and character of growth.

§ 51.3203 Mature.

Mature means that the onion is fairly well cured, and at least fairly firm.

§51.3204 Fairly firm.

Fairly firm means that the onion may yield slightly to moderate pressure but is not appreciably soft or spongy.

$\S 51.3205$ Fairly well shaped.

Fairly well shaped means that the onion shows the characteristic shape, not appreciably three-, four- or five-sided, thick necked or badly pinched.

§51.3206 Wet sunscald.

Wet sunscald means any sunscald which is soft, mushy, sticky or wet.

§ 51.3207 Doubles.

Doubles means onions which have developed more than one distinct bulb joined only at the base.

§ 51.3208 Bottlenecks.

Bottlenecks means onions which have abnormally thick necks with only fairly well developed bulbs.